

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE 3/17/99	3. REPORT TYPE AND DATES COVERED Final Report, 01 Sept. 1993-31 Nov. 1996		
4. TITLE AND SUBTITLE Interfacial Interactions in Microbial Bioadhesion		5. FUNDING NUMBERS N00014-93-1-1317		
6. AUTHOR(S) David C. White, Robert J. Palmer, Cecily A. Flemming				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) The University of Tennessee Center for Environmental Biotechnology 10515 Research Dr., Suite 300 Knoxville, TN 37932-2575		8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research 800 N. Quincy St. Arlington, VA 22217-5000		10. SPONSORING/MONITORING AGENCY REPORT NUMBER		
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Distribution Unlimited		12b. DISTRIBUTION CODE		
13. ABSTRACT (Maximum 200 words) The objective of this research was to characterize the influence of bacterial cell-surface characteristics (exopolymers, surface charges) on adhesion to substrata. Mutants of <i>Pseudomonas aeruginosa</i> with known surface characteristics were used in adhesion tests under laminar flow in aqueous environments. Spectroscopy, microscopy, and fluorescence techniques were used to quantify the relative adhesion propensity of the various mutants. It was discovered that exopolymer is not required in large amounts for adhesion however lipopolysaccharide chemistry seems to play a role in adherence and biofilm accumulation.				
19990323 058				
14. SUBJECT TERMS Bioadhesion, Biofilm, Exopolysaccharide, Microbial			15. NUMBER OF PAGES 3	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT UL	

FINAL REPORT

Grant #: N00014-93-1-1317

R&T Code: 4107-009

PRINCIPAL INVESTIGATOR: Dr. D. C. White, R. J. Palmer, C. A. Flemming

INSTITUTION: Center for Environmental Biotechnology, University of Tennessee at Knoxville

EMAIL: milipids@aol.com

GRANT TITLE: Interfacial Interactions in Microbial Bioadhesion

AWARD PERIOD: 1 September 1993 - 31 August 1996, extended until 31 Nov 1996

OBJECTIVE: To determine the effect of changes in exopolysaccharide structure of bacteria on attachment and biofilm formation.

APPROACH: Utilize mutants of *Pseudomonas aeruginosa* with structural changes in the exopolysaccharides (alginate and lipopolysaccharide) to examine attachment in laminar flow cells by tryptophan fluorescence, direct cell counts and infrared spectroscopy, and to monitor effects of attachment on alginate synthesis by bioluminescence with a *lux-algD* reporter strain. Determine changes in electrostatic and hydrophobic properties of adherence mutants.

ACCOMPLISHMENTS: Attachment of mucoid (wild type; produces exopolymer) and of nonmucoid (mutant; reduced exopolymer production) *P. aeruginosa* cells to germanium prisms was monitored by ATR-FTIR. Both cell types attached and formed biofilms indicating that copious production of exopolymer is not required for attachment or for biofilm formation by these organisms.

Attachment and biofilm formation of *P. aeruginosa* strains O6 (wild type, complete lipopolysaccharide [LPS]), R5 (mutant, lacks almost all LPS) and A28 (mutant, LPS composition intermediate between that of O6 and that of R5) was investigated in laminar flowcells and evaluated by tryptophan fluorescence or by confocal microscopy. Strain R5 attached rapidly and formed densely packed, confluent biofilms of uniform thickness. Strain O6 attached slower and did not grow as rapidly as did strain R5; total biomass in O6 biofilms was lower than for strain R5. Microscopic appearance of the initial biofilms was (in contrast to that of strain R5) patchy with distinct hemispherical microcolonies. Strain A28 did not differ significantly in attachment or biofilm biomass accumulation than strain R5, although the microscopic appearance of the A28 biofilm was intermediate between that of R5 and that of O6.

Hydrophobic Interaction Chromatography and electrostatic Interaction Chromatography were used to investigate the cell-surface characteristics of the *P. aeruginosa* LPS mutants. Strains A28 and R5 were found to be significantly more hydrophobic and higher in surface charge than strain O6. A28 and R5 are known to lack the long-chain LPS subunit ("B-band" LPS).

CONCLUSIONS: Molecular interactions between cells and substrata involved in bacterial adherence and biofilm formation can be investigated nondestructively and in real time. Information on changes in the chemical composition of the biofilms can be acquired by FTIR spectroscopy. This approach can be used to determine environmental effects on exopolymer production and biochemistry. We have shown that exopolymer production appears not to be significant in biofilm formation by these *P. aeruginosa* strains although it certainly plays a role in the physiology of the bacterial community.

In contrast to the exopolymer data, LPS composition does appear to play an important role in attachment, in biofilm formation and in biofilm architecture in the *P. aeruginosa* strains examined. The presence of B-band LPS seems to reduce overall cell-surface charge and hydrophobicity, thereby reducing the propensity for attachment and biofilm growth on the substrata we have investigated (glass, stainless steel).

PUBLICATIONS AND ABSTRACTS

1. Nivens, D.E., J. Schmit, J. Sniateki, T. Anderson, J.Q. Chambers, D.C. White (1993) Multi-channel AFT/FTIR spectrometer for on-line examination of microbial biofilms. *Appl. Spectroscopy* 47:668-671.
2. White, D.C. (1993) *In situ* measurement of microbial biomass, community structure, and nutritional status. *Phil. Trans. Roy. Soc. Lond., Series 344A*:59-67.
3. Franklin, M.J., C.E. Chetan, P. Gacesa, A. Sonesson, D.C. White, D.E. Ohman (1994) *Pseudomonas aeruginosa* AlgG is a processed, polymer level alginate C5-mannuronan epimerase. *J. Bacteriol.* 176:1821-1830.
4. Nivens, D.E., M.E. Franklin, D.C. White, D.E. Ohman (1994) Effect of alginate and acetylation of alginate on formation of biofilms by *Pseudomonas aeruginosa*. Abstract D-41; American Society for Microbiology Annual Meeting; Las Vegas, Nevada
5. Rice, J.F., R.F. Fowler, A.A. Arrage, D.C. White, G.S. Sayler (1994) Effects of external stimuli on environmental bacterial strains harboring an *algD-lux* bioluminescent reporter plasmid for the study of corrosive biofilms. *J. Indust. Microbiol.* 15:318-328.
6. Wallace, W.H., J.F. Rice, D.C. White, G.S. Sayler (1994) Distribution of alginate genes in bacterial isolates from corroded metal surfaces. *Microb. Ecol.* 27: 213-223.
7. Wallace, W.H., J.T. Fleming, D.C. White, G.S. Sayler (1994) An *algD*-bioluminescent reporter plasmid to monitor alginate production in biofilms. *Microb. Ecol.* 27:225-239.
8. Nivens, D.E., R.J. Palmer Jr., D.C. White (1995) Continuous, nondestructive monitoring of microbial biofilms: a review of on-line techniques. *J. Indust. Microbiol.* 15:263-276

9. Palmer, R.J.Jr., D.E. Caldwell (1995) Flowcell Study of Oral Biofilm (Plaque) Communities. Abstract Q-339, American Society for Microbiology Annual Meeting; Washington D.C.
10. Palmer, R.J. Jr., D.E. Caldwell (1995) A flowcell for the study of plaque removal and regrowth. J. Microbiol. Meth. 24:171-182
11. Flemming, C.A., A.A. Arrage, D.C. White (1996) Autofluorescence monitoring of specific bacterial populations. Workshop; American Society for Microbiology conference on Microbial Biofilms. Salt Lake City, Utah
12. Flemming, C.A., K.M. Harter, D.C. White (1996) Determination of cell surface charge and its role in adhesion of *Pseudomonas aeruginosa* LPS mutants. Abstract A-31; American Society for Microbiology conference on Microbial Biofilms; Salt Lake City, Utah
13. Kouznetsova, L.N., T. Kuritz, R.J. Palmer Jr., D.C. White (1996) Effect of chemical modification of glass surfaces on bacterial adhesion. Abstract A-49; American Society for Microbiology conference on Microbial Biofilms; Salt Lake City, Utah
14. Palmer, R.J. Jr., T. Kuritz, L.N. Kouznetsova, D.C. White (1996) Bacterial adhesion to organosiloxane coated surfaces. Oral Presentation; American Society for Microbiology conference on Microbial Biofilms; Salt Lake City, Utah
15. Kuritz, T., L.N. Kouznetsova, C.A. Flemming, R.J. Palmer, Jr., K.M. Harter, D.C. White (1996) Generation and characterization of mutants of *Oceanospirillum* sp. Abstract A-56; American Society for Microbiology conference on Microbial Biofilms; Salt Lake City, Utah
16. Flemming, C.A., R.J. Palmer Jr., D.C. White (1996) Hydrophobicity and attachment of LPS mutants of *Pseudomonas aeruginosa*. Abstract Q-242; American Society for Microbiology Annual Meeting; New Orleans, LA
17. Flemming, C.A., R.J. Palmer Jr., A.A. Arrage, H.C. van der Mei, D.C. White (1998) Cell-surface biochemistry alters biofilm development in *Pseudomonas aeruginosa* lipopolysaccharide mutants. Biofouling 13:213-231